

INTRAVENOUS SAPHNELO® (ANIFROLUMAB) INFUSION CHART

For use by trained healthcare professionals; complete according to institutional policy and physician instructions.

Note*: Saphnelo 300 mg is administered by IV infusion over 30 minutes. It must not be administered by IV push or bolus. Use a sterile, low-protein-binding 0.2–15 micron in-line or add-on filter. If an infusion reaction occurs, the infusion rate may be slowed or interrupted.

Date		Time	
Patient Name		MRN	
Weight (kg)		Dose	300 mg

Saphnelo Infusion & Monitoring

Time from infusion start	Minutes	Vitals			Infusion Reactions					
		B/P	PR	Temp	Headache	Nausea	Vomiting	Fatigue	Dizziness	Others
Start	0									
Mid-infusion	15									
Completion	30									
Post-infusion	45									
Post-infusion	60									

Previous / Prior Infusion Reaction History

Previous Saphnelo infusion reaction?	☐ No ☐ Yes	If yes, date:	
Reaction type / symptoms		Severity:	☐ Mild ☐ Moderate ☐ Severe
Premedication used?	☐ No ☐ Yes	Details:	
Other relevant history		Clinician initials:	

Infusion Reaction / Intervention Record

Time	Signs / Symptoms	Infusion rate / action taken	Treatment / intervention	Outcome	Initials

Additional comments / observations:

Clinician name / signature: _____ Date / time: _____

**Reference note: Administration details are based on current Saphnelo product information.*